

# Neuroimaging

Ritsuko K. Pooh\*

CRIFM Clinical Research Institute of Fetal Medicine PMC, Osaka, Japan

**Abstract:** Transvaginal high-resolution ultrasound and three-dimensional (3D) ultrasound have been establishing sonoembryology in the first trimester as well as neurosonography. Fetal brain is rapidly developing and changing its appearance week by week during pregnancy. The brain is the most important organ but it is quite hard to observe detailed structure of this organ by conventional transabdominal sonography. It is possible to observe the whole brain structure by magnetic resonance imaging in the post half of pregnancy, but it is difficult in the first half of gestation and transvaginal high-resolution 3D ultrasound is the most powerful modality. As for brain vascularization, main arteries and veins have been demonstrated and evaluated in various CNS conditions. Transvaginal high-resolution 3D ultrasound can demonstrate cerebral fine vascular anatomy such as medullary vessels and it is greatly expected to estimate neurological prognosis in relation with vascular development during fetal period.

**Keywords:** Embryo, Fetus, Central nervous system, Transvaginal scan, 3D Ultrasound, Sonoembryology, Sonogenetics, Neuroimaging, Medullary vessels, Neuroangiography.

## INTRODUCTION

Imaging technologies have been remarkably improved and contributed to prenatal evaluation of fetal central nervous system (CNS) development and assessment of CNS abnormalities in utero.

Conventional transabdominal ultrasonography, by which it is possible to observe fetuses through maternal abdominal wall, uterine wall and sometimes placenta, has been most widely utilized for antenatal imaging diagnosis. By transabdominal approach, whole central nervous system of fetuses can be well demonstrated, for instance, the brain in the axial section and the spine in the sagittal section. However, transabdominal approach to the fetal CNS, has several obstacles such as maternal abdominal wall thickness, placenta and fetal cranial bones and it is difficult to obtain clear and detailed images of fetal CNS structure.

Introduction of high-frequency transvaginal transducer has contributed to establishing “sonoembryology”[1] and recent general use of transvaginal sonography in early pregnancy enabled early diagnoses of major fetal anomalies [2,3]. In the middle and late pregnancy, fetal CNS is generally evaluated through maternal abdominal wall. The brain, however, is three-dimensional structure, and should be assessed in basic three planes of sagittal, coronal and axial sections. Sonographic assessment of the fetal brain in the sagittal and coronal sections, requires an approach from fetal parietal direction. Transvaginal sonography of the fetal brain opened a new field in medicine, “neurosonography”[4]. Transvaginal approach to the normal fetal brain during the second and third trimester was introduced in the beginning

of 1990s. It was the first practical application of three-dimensional CNS assessment by two-dimensional (2D) ultrasound [5]. Transvaginal observation of the fetal brain offers sagittal and coronal views of the brain from fetal parietal direction [6-9] through the fontanelles and/or the sagittal suture as ultrasound windows. Serial oblique sections [4] via the same ultrasound window reveal the intracranial morphology in detail. This method has contributed to the prenatal assessment of congenital CNS anomalies and acquired brain damage *in utero*.


The aim of the paper is to introduce the updated transvaginal high-resolution 3D neuroimaging and neuroangiography for precise personal diagnosis of CNS abnormalities.

## TRANSVAGINAL HIGH-RESOLUTION 3D NEUROIMAGING

Three-dimensional (3D) ultrasound is one of the most attractive modalities in the field of fetal ultrasound imaging. Automatic scan by dedicated 3D transducer produces motor driven automatic sweeping and is called as a fan scan. With this method, shifting and/or changing angle of the transducer is not required during scanning and scan duration lasts only several seconds. After acquisition of the target organ, multiplanar imaging analysis and tomographic imaging analysis are possible. Combination of both transvaginal high-resolution sonography and 3D ultrasound [10-17] may be a great diagnostic tool for evaluation of three-dimensional structure of fetal CNS. Recent advanced 3D ultrasound equipments have several useful functions such as:

- Surface anatomy imaging of spinal development, cranial surface and brain surface.
- Bony structural imaging of the calvaria and vertebrae.
- Multiplanar imaging of the intracranial structure.

\*Address correspondence to this author at the CRIFM Clinical Research Institute of Fetal Medicine PMC 7-3-7, Uehommachi, Tennoji, Osaka #543-0001, Japan; Tel +81-6-6775-8111; Fax: +81-6-6775-8122; E-mail: evp-pooh@fetal-medicine-pooh.jp



**Fig. (1). Cephalocele at 12 weeks of gestation.**

Left figure; Tomographic ultrasound imaging (TUI). Right figure ; 3D reconstructed image.



**Fig. (2). Craniovertebral imaging in an acranial fetus.**

Left upper image is 2D ultrasound imaging and left lower4 image is multiplanar imaging. Right picture shows 3D bony structure of cranio-vertebral part. Acrania and opening of cervical vertebrae are demonstrated.

- Tomographic ultrasound imaging of fetal brain in any cutting section.
- Thick slice imaging of the intracranial structures (Volume contrast imaging, VCI).
- Simultaneous volume contrast imaging of the same section or vertical section of fetal brain structure.
- Volume calculation of target organs such as intracranial cavity, ventricle, choroid plexus and intracranial lesions.


- Inversion mode demonstrating hypochoic parts such as ventricles and cystic lesions.

It is well known that 3D ultrasound demonstrates the surface anatomy. Depiction of CNS and facial abnormalities as well as anomalies of the extremities is often complicated. Therefore, surface reconstructed images are helpful. In early pregnancy, small cephalocele can be demonstrated (Fig.1). Bony structural imaging of the calvaria and vertebrae are useful in cases of craniosynostosis and spina bifida. Fig. (2) shows the craniovertebral imaging in an acranial fetus. Sur-




**Fig. (3). Normal brain 3D surface anatomy in 19 and 30 weeks of gestation.**

Notice obviously different appearance of the smooth brain surface at 19 weeks and matured brain surface with gyration at 30 weeks.



**Fig. (4). Abnormal brain 3D surface anatomy in pachygyria.**

The brain surface with less gyration at 33 weeks of gestation is well demonstrated in case of pachygyria.




**Fig. (5). Tomographic ultrasound imaging of normal brain in coronal cutting section at 20 weeks.**


face anatomy imaging can demonstrate the brain surface shown in the Fig. (3 and 4). Migration occurs during 3 to 5 gestational month but the gyration and sulcation can be visualized in the late pregnancy. Fig. (3) shows obviously different appearance of the brain surface at 19 and 30 weeks of gestation. Fig. (4) shows the brain surface with less gyration at 33 weeks of gestation seen in a case of pachygyria. Thus,

brain development can be objectively understandable by surface reconstructed imaging.

In multiplanar imaging of the brain structure and tomographic ultrasound imaging (TUI, Fig. (1) left and Fig. (5) are quite helpful to understand intracranial detailed brain structure. It is possible to compare the anatomy in ex-



**Fig. (6).** Changing appearance of Sylvian fissure during the second and early third trimesters




**Fig. (7).** Early fetal intracranial hemorrhage/porencephaly case.

(Upper figures) Dedicated transvaginal high-frequency sonography detected the subtle change of ventricular wall between 16 and 17 weeks of gestation (arrows). (Left lower) Ventricular wall is irregular and hyperechoic (red arrows), indicating intraventricular hemorrhage. (Right lower) MR image of the same fetus at 20 weeks. Asterisks indicate the missing brain (porencephaly) because of early brain hemorrhage.

actly same cutting sections. Fig. (6) shows the changing appearance of Sylvian fissure during the second and early third trimesters. Fig. (7) shows early intracranial hemorrhage case and dedicated transvaginal high-frequency sonography detected the subtle change of ventricular wall between 16 and 17 weeks of gestation. TUI demonstrates multi-parallel cutting sections and is quite similar imaging technology to magnetic resonance imaging. The superior point of TUI to MRI is that it is easily possible to change slice width, to rotate the images, to magnify images, and to rotate images to any directions. This function is extremely useful for detailed CNS assessment and also for consultation to neurosurgeons and neurologists. Fig. (8) shows abnormal migration disorder in the second trimester. The intracranial brain structure is quite abnormal and it is difficult to obtain the orientation but TUI is quite helpful to show the intracranial structure very clearly. Thick slice imaging of the intracranial structure and simultaneous volume contrast imaging (VCI) of the same plane or vertical plane of conventional 2D image are often convenient to observe the gyral formation and inside lateral


ventricles [14]. The premature brain image obtained by use of VCI clearly demonstrates anatomical CNS structure.

Volume extracted image and volume calculation of the fetal brain have been reported from 1990s [18-22]. On three orthogonal images, the target organ can be traced automatically or manually with rotation of volume imaging data. After tracing, volume extracted image is demonstrated and data of volume calculation are shown. 3D fetal brain volume measurements have a good intraobserver and interobserver reliability [23, 24] and could be used to determine estimated gestational age. Volume analysis by 3D ultrasound provides exceedingly informative imaging data. Volume analysis of the structure of interest provides an intelligible evaluation of the brain structure in total, and longitudinal and objective assessment of enlarged ventricles and intracranial occupying lesions (Fig. 9). Any intracranial organ can be chosen as a target for volumetry no matter how distorted its shape and appearance may be. In new method of inversion mode, the cystic portions within the volume are displayed in their entirety as an echogenic area, while the grayscale portions of




**Fig. (8). Abnormal migration disorder in the second trimester.**

(Upper figures) TUI in the sagittal (left) and coronal (right). The intracranial brain structure is chaotic and it is difficult to obtain the orientation but TUI is quite helpful to show the intracranial structure very clearly. Lower left picture shows the brain surface at autopsy in the same case. Abnormal gyration which cannot exist at 21 weeks is seen. Lower right figure is normal smooth brain surface at the same gestation for comparison.



**Fig. (9). 3D volume extraction and volumetric analysis of lateral ventricle and intracranial cavity.**

Each volume of right (RV) and left ventricles (LV) and intracranial cavity volume can be calculated by 3D volumetry. Total ventricular volume / Intracranial cavity (ICC) volume shows ventricles occupying rate and it is useful for longitudinal assessment of fetal ventriculomegaly.



**Fig. (10). TUI and inversion mode of enlarged ventricle at 19 weeks of gestation.**  
Ventricular appearance is objectively demonstrable by inversion mode.

the image are rendered as transparent [25] and recently it has been applied in fetal diagnosis [26, 27].


Inversion mode is inverted imaging technology which shows volume of only hypoechoic parts. Fig. (10) shows TUI and inversion-mode images of enlarged ventricles seen at 19 weeks of gestation.

Thus, owing to transvaginal high-resolution 3D neuroimaging, details of CNS normal development and maldevelopment have been elucidated.


#### **TRANSVAGINAL HIGH-RESOLUTION 3D NEUROANGIOGRAPHY**

The author first reported brain circulation demonstrated by Transvaginal 2D power Doppler in 1996 [28]. Thereafter, transvaginal 3D power Doppler assessment of fetal brain


vascularity was successful [21, 29]. Recently, owing to the advanced technology of directional power Doppler, furthermore sophisticated 3D angiostructural image have been able to be demonstrated with vascular direction [30]. In the first trimester, it is possible to demonstrate the fetal cranial circulation clearly as early as 7 weeks of gestation shown in Fig. (11). Brain circulation can be well documented during the first trimester. Fig. (12) shows the fetal brain circulation at 14 weeks by 3D bidirectional power Doppler. Furthermore, recent high-frequent transvaginal neuroscan has been able to demonstrate the medullary vessels from the cortex towards subependymal area. Medullary veins are demonstrated between pia mater and longitudinal vein of Schlessinger in the anterior coronal cutting section and between brain surface and periventricular subependymal area in the



**Fig. (11). Early cranial circulation at 7 weeks of gestation.** Left upper shows B-mode image of sagittal section and right figure shows fetal circulation by bidirectional power Doppler technology.



**Fig. (12). Brain circulation at 14 weeks by 3D bidirectional power Doppler.** (left) Volume contrast imaging in coronal cutting section. (middle) 3D bidirectional power Doppler reconstructed image with B-mode. (right) Brain angiography by 3D bidirectional power Doppler. Those 3 images are obtained by a single acquisition.



**Fig. (13). Medullary vessels in normal 29-week fetus by 3D bidirectional power Doppler.**

(upper) Coronal cutting section of 2D (left) and 3D reconstructed images (right).

(lower) Parasagittal cutting section of 2D (left) and 3D reconstructed images (right).

Medullary veins are demonstrated between pia mater and longitudinal vein of Schlesinger in the anterior coronal cutting section and between brain surface and periventricular subependymal area in the parasagittal section.

parasagittal section [30] (Fig. 13). Medullary vessels are detectable from early second trimester and they are rapidly developing during second trimester. Medullary vessels are developing according to advancing gestational age in normal fetuses and maldevelopment of those vessels may relate with the adverse postnatal neurological prognosis [30].

#### FUTURE ASPECT

As described in this article, transvaginal high-resolution ultrasound technology has greatly contributed to fetal neuroscience. Neurological prognosis should be longitudinally and carefully evaluated according to precise diagnoses. Considering future development in the field of fetal neurology, those precise morphological detection by transvaginal high-resolution neuroimaging should be combined with four-dimensional ultrasound research on fetal behavior [31, 32] and molecular genetics which has recently been remarkably contributed to prenatal diagnosis as *Sonogenetics* [33].

#### ACKNOWLEDGEMENT

None declared.

#### CONFLICT OF INTEREST

None declared.

#### REFERENCES

- [1] Timor-Tritsch IE, Peisner, DB, Raju S. Sonoembryology: an organ-oriented approach using a high-frequency vaginal probe. *J Clin Ultrasound* 1990; 18: pp. 286-98.
- [2] Pooh RK. B-mode and doppler studies of the abnormal fetus in the first trimester. In: Chervenak FA, Kurjak A, Eds. *Fetal medicine*. Carnforth: Parthenon Publishing 1999; pp. 46-51.
- [3] Pooh RK, Shiota K, Kurjak A. Imaging of the human embryo with magnetic resonance imaging microscopy and high-resolution transvaginal 3-dimensional sonography: human embryology in the 21<sup>st</sup> century. *Am J Obstet Gynecol* 2011; 204(1): pp. 77.e1-16.
- [4] Timor-Tritsch IE, Monteagudo A. Transvaginal fetal neurosonography: standardization of the planes and sections by anatomic landmarks. *Ultrasound Obstet Gynecol* 1996; 8: pp. 42-7.
- [5] Monteagudo A, Reuss ML, Timor-Tritsch IE. Imaging the fetal brain in the second and third trimesters using transvaginal sonography. *Obstet Gynecol* 1991; 77: pp. 27-32.
- [6] Monteagudo A, Timor-Tritsch IE, Moomjy M. In utero detection of ventriculomegaly during the second and third trimesters by transvaginal sonography. *Ultrasound Obstet Gynecol* 1994; 4: pp. 193-8.
- [7] Pooh RK, Nakagawa Y, Nagamachi N, *et al.* Transvaginal sonography of the fetal brain: detection of abnormal morphology and circulation. *Croat Med J* 1998; 39: pp. 147-57.
- [8] Pooh RK, Maeda K, Pooh KH, Kurjak A. Sonographic assessment of the fetal brain morphology. *Prenat Neonatal Med* 1999; 4: pp. 18-38.
- [9] Pooh RK. Three-dimensional ultrasound of the fetal brain. In: Kurjak A, Ed. *Clinical application of 3D ultrasonography*. Carnforth: Parthenon Publishing 2000; pp. 176-80.
- [10] Pooh RK, Pooh KH, Nakagawa Y, Nishida S, Ohno Y. Clinical application of three-dimensional ultrasound in fetal brain assessment. *Croat Med J* 2000; 41: pp. 245-51.
- [11] Timor-Tritsch IE, Monteagudo A, Mayberry P. Three-dimensional ultrasound evaluation of the fetal brain: the three horn view. *Ultrasound Obstet Gynecol* 2000; 16: pp. 302-6.
- [12] Monteagudo A, Timor-Tritsch IE, Mayberry P. Three-dimensional transvaginal neurosonography of the fetal brain: 'navigating' in the volume scan. *Ultrasound Obstet Gynecol* 2000; 16: pp. 307-13.
- [13] Pooh RK, Nagao Y, Pooh KH. Fetal neuroimaging by transvaginal 3D ultrasound and MRI. *Ultrasound Rev Obstet Gynecol* 2006; 6: pp. 123-34.
- [14] Pooh RK, Kurjak A. 3D and 4D sonography and magnetic resonance in the assessment of normal and abnormal CNS demelop-

- ment: alternative or complementary. *J Perinat Med* 2011; 39(1): pp. 3-13.
- [15] Pooh RK, Pooh KH. Fetal neuroimaging with new technology. *Ultrasound Rev Obstet Gynecol* 2002; 2: pp. 178-81.
- [16] Pooh RK, Maeda K, Pooh KH. An atlas of fetal central nervous system disease. diagnosis and management. London, New York: Parthenon CRC Press 2003.
- [17] Pooh RK. Neuroanatomy Visualized by 2D and 3D. In: Pooh RK, Kurjak A, Eds. *Fetal neurology*. New Delhi: Jaypee Brothers Medical Publishers 2009; pp. 15-38.
- [18] Blaas HG, Eik-Nes SH, Kiserud T, Berg S, Angelsen B, Olstad B. Three-dimensional imaging of the brain cavities in human embryos. *Ultrasound Obstet Gynecol* 1995; 5: pp. 228-32.
- [19] Blaas HG, Eik-Nes SH, Berg S, Torp H. *In-vivo* three-dimensional ultrasound reconstructions of embryos and early fetuses. *Lancet* 1998; 352: pp. 1182-6.
- [20] Pooh RK. Fetal Brain Assessment by Three-Dimensional Ultrasound. In: Kurjak A, Kupesic S, Eds. *Clinical application Of 3D sonography*. Carnforth, UK, Parthenon Publishing 2000; pp. 171-9.
- [21] Pooh RK, Pooh KH. Transvaginal 3D and Doppler ultrasonography of the fetal brain. *Semin Perinatol* 2001; 25: pp. 38-43.
- [22] Pooh RK, Pooh KH. The assessment of fetal brain morphology and circulation by transvaginal 3D sonography and power Doppler. *J Perinat Med* 2002; 30: pp. 48-56.
- [23] Endres LK, Cohen L. Reliability and validity of three-dimensional fetal brain volumes. *J Ultrasound Med* 2001; 20: pp. 1265-9.
- [24] Roelfsema NM, Hop WC, Boito SM, Wladimiroff JW. Three-dimensional sonographic measurement of normal fetal brain volume during the second half of pregnancy. *Am J Obstet Gynecol* 2004; 190(1): pp. 275-80.
- [25] Benacerraf BR. Inversion mode display of 3D sonography: applications in obstetric and gynecologic imaging. *Am J Roentgenol* 2006; 187(4): pp. 965-71.
- [26] Kusanovic JP, Nien JK, Gonçalves LF, *et al*. The use of inversion mode and 3D manual segmentation in volume measurement of fetal fluid-filled structures: comparison with Virtual Organ Computer-aided AnaLysis (VOCAL). *Ultrasound Obstet Gynecol* 2008; 31(2): pp. 177-86.
- [27] Hata T, Dai SY, Kanenishi K, Tanaka H. Three-dimensional volume-rendered imaging of embryonic brain vesicles using inversion mode. *J Obstet Gynaecol Res* 2009; 35(2): pp. 258-61.
- [28] Pooh RK, Aono T. Transvaginal power Doppler angiography of the fetal brain. *Ultrasound Obstet Gynecol* 1996; 8: pp. 417-21.
- [29] Pooh RK. Two-dimensional and three-dimensional Doppler angiography in fetal brain circulation. In: Kurjak A, Ed. *3D power doppler in obstetrics and gynecology*. Carnforth: Parthenon Publishing 1999; pp. 105-11.
- [30] Pooh RK, Kurjak A, Tikvica A. Normal and abnormal brain vascularity. In: Pooh RK, Kurjak A, Eds. *Fetal neurology*. New Delhi: Jaypee Brothers Medical Publishers 2009; pp. 39-58.
- [31] Amiel-Tison C, Gosselin J, Kurjak A. Neurosonography in the second half of fetal life: a neonatologist's point of view. *J Perinat Med* 2006; 34(6): pp. 437-46.
- [32] Kurjak A, Abo-Yaqoub S, Stanojevic M, *et al*. The potential of 4D sonography in the assessment of fetal neurobehavior--multicentric study in high-risk pregnancies. *J Perinat Med* 2010; 38(1): pp. 77-82.
- [33] Pooh RK, Choy KW, Leung TY, Lau TK. Sonogenetics-A breakthrough in prenatal diagnosis. *Donald Sch J Ultrasound Obstet Gynecol* 2011; 5(1): pp. 75-9.

Received: January 22, 2012

Revised: March 03, 2012

Accepted: April 05, 2012

© Ritsuko K. Pooh; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.